



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Spontaneously occurring melanoma in animals and their relevance to human melanoma

Citation for published version:

van der Weyden, L, Brenn, T, Patton, EE, Wood, GA & Adams, DJ 2020, 'Spontaneously occurring melanoma in animals and their relevance to human melanoma', *The Journal of Pathology*.
<https://doi.org/10.1002/path.5505>

Digital Object Identifier (DOI):

[10.1002/path.5505](https://doi.org/10.1002/path.5505)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

The Journal of Pathology

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.





Spontaneously occurring melanoma in animals and their relevance to human melanoma

Journal:	<i>The Journal of Pathology</i>
Manuscript ID	20-348.R1
Wiley - Manuscript type:	Review
Date Submitted by the Author:	n/a
Complete List of Authors:	van der Weyden, Louise; Sanger Institute, Experimental Cancer Genetics (T113) Brenn, Thomas; University of Calgary, Arnie Charbonneau Cancer Institute Patton, Elizabeth; MRC , Human Genetics Unit Geoffrey, Geoffrey; Ontario Veterinary College, Department of Pathobiology Adams, David; Wellcome Trust Sanger Institute, Experimental Cancer Genetics
Portal Keywords:	Animal model, Dermis, DNA sequencing, Epidermis, In vivo models, Neoplasia, Skin
Tissue:	
Pathology:	
Technique:	

SCHOLARONE™
Manuscripts

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

Spontaneously occurring melanoma in animals and their relevance to human melanoma

8

9

Running title: Spontaneously occurring melanoma in animals

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

Louise van der Weyden^{1*}, Thomas Brenn², E. Elizabeth Patton³, Geoffrey A Wood⁴, David J. Adams¹

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

¹Wellcome Sanger Institute, Wellcome Genome Campus, Hinxton, Cambridge, CB10 1SA, UK.

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

²Arnie Charbonneau Cancer Institute, University of Calgary, Calgary, Alberta, T2N 1N4, Canada.

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

³MRC Human Genetics Unit, The MRC Institute of Genetics and Molecular Medicine, The University of Edinburgh, Western General Hospital, Crewe Road, Edinburgh, EH4 2XU, UK.

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

⁴Ontario Veterinary College, University of Guelph, Guelph, N1G 2W1, Canada.

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

*Corresponding author:

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

Louise van der Weyden

Experimental Cancer Genetics

Wellcome Sanger Institute

Wellcome Genome Campus, Hinxton, Cambridge, CB10 1SA, United Kingdom

Tel: +44-1223-834-244

Email: lvdw@sanger.ac.uk

56

57

58

59

60

Conflict of interest statement: The authors declare there are no conflicts of interest.

56

57

58

59

60

Word count for main text: 7,118

Abstract

In contrast to other cancer types, melanoma incidence has been increasing over the last 50 years, and while it still represents less than 5% of all cutaneous malignancies, melanoma accounts for the majority of skin cancer deaths, due to its propensity to metastasize. Whilst melanoma most commonly affects the skin, it can also arise in mucosal surfaces, the eye, and the brain. In order for new therapies to be developed, a better understanding of the genetic landscape, signalling pathways and tumour-microenvironmental interactions are needed. This is where animal models are of critical importance. The mouse is the foremost used model of human melanoma. Arguably this is due to its plethora of benefits as a laboratory animal, however, it is important to note that unlike humans, melanocytes are not present at the dermal-epidermal junction in mice and mice do not develop melanoma without genetic manipulation. In contrast, there are numerous case reports of animals that spontaneously develop melanoma, ranging from sharks and parrots to hippos and monkeys. In addition, there are also several domesticated and laboratory-bred animals that spontaneously develop melanoma or UV-induced melanoma, specifically, fish, opossums, pigs, horses, cats and dogs. In this review, we look at spontaneously occurring animal 'models' of melanoma and discuss their relevance to the different types of melanoma found in humans.

Keywords

Melanoma, naturally occurring, UV-induced, animal model, fish, pig, cat, horse, dog.

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

1. Introduction

Melanoma is a tumour that arises from uncontrolled proliferation of melanocytes (pigment-producing cells). Although the most common form of melanoma is cutaneous, it can also arise from melanocytes in the mucosal surfaces (mucosal melanoma), the eye (ocular melanoma), and leptomeninges (leptomeningeal melanoma). Although all subtypes of melanomas are derived from melanocytes and thus share the same embryonic origin and cellular function, the etiopathogenesis and biological behaviour of these melanoma subtypes are very different with distinct landscapes of genetic alterations and different metastatic routes. The major signalling pathways implicated in melanoma are shown in Figure 1 and the most commonly mutated genes for each melanoma subtype are shown in Table 1.

1. 1

Melanoma developing on sun-exposed skin

The current 4th edition of the World Health Organization (WHO) classification of skin tumours separates melanocytic tumours according to sun-exposure: melanoma arising in intermittently sun-exposed skin, melanoma arising in chronically sun-exposed skin and melanoma arising on sun-protected skin [1]. Chronic and intermittent ultraviolet (UV) light exposures are a major cause of melanoma development in sun-exposed skin, and UV-induced melanoma is the most common type of melanoma in pale/fair skin types (people of European descent), typically occurring at 30-60 years of age (or 70 years of age in cumulative sun-induced damage cases) [2]. The degree of cumulative sun-exposure also influences the histological subtypes of melanoma. Melanomas on chronically sun-exposed skin are typically of the lentigo maligna or desmoplastic variants, while melanoma on intermittently sun-exposed skin is most frequently of the superficial spreading or nodular subtypes. Although cutaneous melanoma represents <5% of all skin malignancies, it accounts for the majority of skin cancer deaths [3], due to its aggressive nature and propensity to metastasise. Whilst the prognosis for

patients with advanced stage melanoma has undoubtedly improved in recent years, with a current five year survival of 52% and median survival of >60 months for those treated with combination immunotherapy (Ipilimumab/Nivolumab) [4], more progress still needs to be made.

1.2 Melanoma developing on sun-protected areas

The large group of melanomas developing on sun-protected skin includes acral melanoma, genital and mucosal melanoma, ocular melanoma and leptomeningeal melanoma. Other rare subtypes of melanoma that arise independent of UV light exposure are Spitz melanoma, melanoma arising in blue naevi and melanoma arising in congenital naevi.

1.2.1 Acral melanoma

Acral melanoma arises on sun-protected sites, including the palms of the hands, soles of the feet, and the nail apparatus (subungal), of the middle-aged and elderly with a mean of around 60 years of age. Although the absolute incidence is independent of ethnicity the relative incidence is highest individuals with darker skin types (people of African, Middle Eastern or Asian ethnicity), typically occurring at 60 years of age [2]. The majority of melanomas are of the acral lentiginous or nodular subtype. Acral melanoma carries a particularly poor prognosis, as it is often detected at a more advanced stage.

1.2.2 Genital and mucosal melanoma

In comparison to cutaneous melanoma, genital and mucosal melanomas (MM) are much rarer, representing only 0.8-3.7% of all melanomas [5]. They are relatively more common in the Asian and black populations where they account for 9-22% of all melanoma cases [5]. This is due to the rarity of UV light associated melanoma subtypes in this patient population. Like acral melanomas, genital and mucosal melanomas typically occur at 60 years of age [2]. The majority of MMs present in the nasal cavity and the accessory sinuses, the oral cavity,

1 anorectum, vulva and vagina, but they can arise in almost every part of mucosal membranes,
2
3 including the parotid glands, oesophagus and middle ear [6]. Histologically, they are most
4
5 commonly of the lentiginous mucosal subtype followed by nodular melanoma. MMs are known
6
7 to behave more aggressively with less favourable prognosis compared to other melanoma
8
9 subtypes. Most MMs occur in occult sites, which together with the lack of early and specific
10
11 signs, contribute to late diagnosis and poor prognosis, with a 5 year survival rate of 25-33%,
12
13 according to disease stage and location [7]; metastasis to lungs and liver occur in 50% of cases
14
15 with head and neck melanoma [7].
16
17
18
19
20
21

22 1.2.3 Ocular melanoma

23
24 Ocular melanoma (OM) is the most common primary cancer affecting the eye, and is
25
26 classified based on the anatomic site of origin: conjunctival melanoma or uveal melanoma
27
28 (UM). The majority of OM originates from the uvea (95%), comprising the posterior uvea
29
30 (choroid 90% and ciliary body 5%) and the anterior uvea (iris 5%). There is a wide age
31
32 distribution for the development of UM, with an average age of 60 years [2]. Although the
33
34 primary tumour can be effectively surgically removed, half of all patients develop metastasis
35
36 (typically the liver or lung) and these patients have a poor prognosis, with a median survival
37
38 time of 4-15 months [8]. Conjunctival melanoma generally presents as a pigmented nodular
39
40 lesion typically on the bulbar conjunctiva and often involves the limbus [9]. Approximately 20-
41
42 30% of conjunctival patients develop metastatic disease, either directly extending into the
43
44 eyeball and orbit or more distant metastases to the lungs, brain, liver, skin, bones and
45
46 gastrointestinal tract [10].
47
48
49
50
51
52
53

54 1.2.4 Primary melanocytic neoplasms of the central nervous system

55
56 Primary melanocytic neoplasms of the central nervous system (CNS), which can be
57
58 divided into either benign or malignant and localised or diffuse, are presumed to arise from the
59
60 melanocytes in the leptomeningeal tissue [11]. However, primary melanoma of the CNS is rare,

with only an incidence of 0.005 per million people [12] (compared to 2.8-3.1 per 100,000 people for cutaneous melanoma [13]), whereas secondary intracranial melanomas are more common, with ~10-40% of melanomas metastasizing to the CNS [14]. Meningeal melanocytomas are solitary, low-grade tumours with no invasion of surrounding structure, with a reported annual incidence of 1 case per 10 million population [15]. In contrast, meningeal melanomas are solitary, highly aggressive, radioresistant tumours that can metastasize, resulting in poor prognosis, with a reported annual incidence of 0.5 cases per 10 million population [16]. The diffuse presenting tumours, generally involving large expanses of the subarachnoid space, are meningeal melanocytosis, which is a proliferation of cytologically bland melanocytic cells, and meningeal melanomatosis, which is a primary melanoma of the CNS. Diffuse meningeal melanocytic tumours are rare and as such population-based incidence is not available, however, they are strongly linked with neurocutaneous melanocytosis, which is a rare disease that is presents before 2 years of age and is typically associated with giant congenital naevi [17]. The prognosis for diffuse meningeal melanocytic tumours, in the setting of neurocutaneous melanocytosis, is extremely poor, with no known effective treatment options and no reports of survivors in the literature [18].

1. 3 Melanoma of unknown primary

Melanoma of unknown primary (MUP) refers to metastatic melanoma occurring in lymph nodes, subcutaneous tissue, or visceral sites in the absence of a detectable primary tumour [19] and accounts for ~2–9% of cases of metastatic melanoma [20]. The most common clinical presentation of MUP is lymph node disease without clinical or radiological evidence of visceral involvement [19] and typically presents in the fifth and sixth decades of life [21]. The origin of MUP is not fully understood and explanations have included regression of primary cells at the site of origin (regression is found in melanoma with a frequency ranging from 10% to 35% [22]), misdiagnosis of the original primary cutaneous melanoma and de novo malignant

transformation of ectopic melanocytes in lymph nodes or other organs [19]. Interestingly, genomic analysis of MUPs has found they have a mutational profile more similar to cutaneous melanoma than that of mucosal melanoma or CNS melanoma, including mutations in *BRAF* and *NRAS* [23], suggesting that regression or misdiagnosis of a previous cutaneous lesion was the primary melanoma.

1. 4 The use of the mouse to model melanoma

In order to study basic mechanisms and pre-clinically test novel therapies, accurate animal models are required. The most popular animal model for melanoma research is the laboratory mouse, and there are numerous mouse models of cutaneous melanoma [reviewed in 24] and uveal melanoma [reviewed in 25], as well as a few models of mucosal melanoma [26] and leptomeningeal melanoma [27]. Although each model has its own distinct advantages and disadvantages (transgenic models, gene knockout/knockin models, xenograft models, or UV/carcinogen-induced models), our understanding of the molecular mechanisms of melanoma development in humans has been greatly improved by modelling various pathways in these mice [28-29]. However, it is worth mentioning that interpreting results from melanoma studies in mice need to take into consideration several key aspects: (i) the heterogeneity of melanoma in humans, which would not be recapitulated in inbred laboratory mice, (ii) the spontaneous development of melanoma in humans, as melanoma does not spontaneously occur in the mouse, and (iii) the differences in the structure of the skin between humans and mice, as melanocytes are located in the epidermis in humans and in the dermis in mice [30], meaning there is a different melanocyte microenvironment, which may account for differences in melanoma development and progression between the two species. Thus it is worth looking at animals that spontaneously develop melanoma (of all subtypes) and assessing what relevance they have to human melanoma as alternative animal models.

2. Melanoma in the 'wild'

There have been reports of melanoma spontaneously occurring in a wide range of species, including fish, reptiles, birds, and mammals, with some examples detailed in Table 2. In reptiles, tumours of pigment-producing cells of the skin are called chromatophoromas, and are subclassified on the basis of the type of pigment, specifically melanophoromas or iridophoromas. A review of chromatophoromas that arose in 26 reptiles (6 snakes, 19 lizards and 1 tortoise) found that 6 of the 20 melanophoromas, were classified as malignant due to the presence of intravascular tumour cells, visceral metastases, high pleomorphism and/or mitotic figures [44]. Microscopically, most of the tumours were composed of spindle cells with varying pigmentation and both melan A and S100 were expressed by all of the tumours (by immunohistochemistry) [44]. Six of the 9 reptiles that were euthanized, either immediately after diagnosis or following recurrence of their tumour, had visceral metastases [44]. Figure 2 shows a metastatic melanophoroma found at autopsy of a 60-year old red-eared slider turtle (*Trachemys scripta elegans*).

However, whilst it is of interest to see the similarities of the histological descriptions of these melanomas with that seen in human melanoma, these case reports demonstrate the low frequency of the spontaneous development of this tumour type in these animals. Indeed, malignant melanomas are rare in fish and extremely rare in reptilian and avian species. In addition, the specialised habitat required for many of these animals, makes them unfeasible for use as a model organism. However, there are several non-genetically-modified laboratory-bred animals that develop melanoma (due to selective breeding of affected individuals), some spontaneously and others after exposure to radiation such as specific species of fish, opossums and pigs. In addition, there several domesticated animals in which the spontaneous development of melanomas is relatively common, such as the dog and horse. These animals offer many important advantages as 'models' of melanoma in humans, which are discussed below.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

3. Laboratory-bred animal models of spontaneously occurring melanoma

3. 1 Fish

The first fish models of melanoma were established in *Xiphophorus* (commonly known as platyfish or swordtails), which are euryhaline and freshwater fish native to Mexico and northern Central America. The *Xiphophorus* model (also known as “Gordon–Kosswig–Anders” melanoma) was originally introduced in the late 1920s and uses interspecies platyfish that spontaneously develop melanoma, in a genetically controlled manner (reviewed in [45]). The platyfish (*X. maculatus*) parental line carries the *spotted dorsal* (*Sd*) macromelanophore pigmentation pattern. These spots are non-malignant melanocytic lesions or hyperpigmentation on the dorsal fin, and the macromelanophore pattern is considered a benign precursor to melanoma. This is controlled by the pigment-cell-specific oncogene locus *Tu* (for tumour), the oncogenic activity of which is suppressed by the tumour suppressor locus *R* (for regulator). In the classical cross, platyfish, which carry *Tu* and *R*, are crossed with swordtails (*X. hellerii*), which carry neither locus and thus have no pigmentation pattern. When the resulting F1 hybrids are backcrossed with swordtails, 25% of progeny inherit the *Tu* oncogene as heterozygotes, but not the *R* locus, and these *Tu*⁻ hybrids develop spontaneous, highly malignant melanoma (Figure 3).

Positional cloning identified the melanoma-inducing *Tu* locus as a novel receptor tyrosine kinase termed *Xmrk* (*Xiphophorus melanoma receptor tyrosine kinase*) [46], that originated from a tandem gene duplication of the *Xiphophorus EGFR* (*epidermal growth factor receptor*) gene, and is the fish orthologue of human *EGFR*. The oncogenic activity of *Xmrk* depends on its overexpression, which correlates with the aggressiveness of the melanoma, as *Xmrk* expression increases with malignancy [47]. *Xmrk* carries two activating mutations that result in a constitutively active receptor whose signals elicit a variety of cellular responses

including proliferation, anti-apoptosis and cell motility, that ultimately result in melanoma development (reviewed in [45]). Interestingly, despite being oncogenic *Xmrk* has been maintained for several million years in the *Xiphophorus* melanoma model, and clues for the positive selection were found in experiments with *Xiphophorus cortezi* that showed females prefer males with an enhanced Sc pattern [48], *Xmrk* males (regardless of Sc phenotype) were more aggressive than *Xmrk*-deficient (wildtype) males [49] and collections of males from four natural populations of these fish found that those with malignant melanomas were significantly larger than both *Xmrk* males without melanomas and *Xmrk*-deficient males [50]. Based on genetic linkage analyses the *Cdkn2ab* gene was proposed as the critical component of the tumour suppressor *R* locus (so named due to 55% amino acid homology with the human tumour suppressor genes *CDKN2A* and *CDKN2B* that are frequently mutated in melanoma) [51]. *Cdkn2ab* has been subsequently shown to have a melanoma-suppressive function in *Xmrk* transgenic medaka models [47], but it is not clearly established if the *R* locus is *Cdkn2ab*.

The relevance of the *Xiphophorus* model to human melanoma stems from the fact that *Xmrk*, being an oncogenic version of the well-studied EGFR, uses several signalling pathways already known for the EGFR, including the Ras/Raf/MAPK (mitogen-activated protein kinase) pathway that is commonly mutated in human melanomas (reviewed in [52]). This fish model significantly contributed to the understanding of the importance of the receptor tyrosine kinase (RTK)–Ras–MAPK pathway signalling in melanoma, along with engineered melanoma models in mice, cell line studies and human chromosomal rearrangements (reviewed in [53]). Similarly, *Xmrk* also induces the PI3K signalling pathway, and it was the discovery of the constitutive activation of STAT5 correlating with the aggressiveness of melanoma in *Xiphophorus* [54], that prompted investigation of the role of STAT5 human melanoma [55].

Xiphophorus also represent a relevant model as they are one of only a few animal models for which the induction of melanomas can occur by exposure to UV radiation alone and studies in these fish were important for proving the role that solar UV-B radiation plays in the formation of cutaneous melanoma [56]. More recently, gene expression profiling of melanomas

1 produced in these fish showed the high-expressing *Xmrk Xiphophorus* melanomas share
2 transcriptomic features and molecular functions of highly proliferative, dedifferentiated human
3 melanoma [57].
4
5
6
7
8
9

10 **3.2 Opossum**
11

12 The gray short-tailed opossum, *Monodelphis domestica* is a small (~100 g) pouchless
13 marsupial native to South America that has been maintained in outbred colonies in laboratories
14 since 1978 (reviewed in [58]). *Monodelphis* has the distinction of being the only naturally
15 existing mammal, other than humans, known to be susceptible to melanoma in response to UV
16 radiation alone. It was long suspected that UV radiation was involved in the etiology of
17 cutaneous melanoma in humans, however, it was studies in *Monodelphis* that proved that UV
18 radiation can act as a complete carcinogen for melanoma induction, and that DNA damage is
19 involved in melanoma formation [59]. UV (primarily UV-B)-exposed *Monodelphis* develop a
20 variety of hyperplastic and neoplastic skin lesions on their shaved back, including malignant
21 melanoma [60] and UV exposure of neonatal *Monodelphis* results in the development of
22 naevus with further repeated UV exposure into adulthood (up to 45 weeks of age) resulting in
23 progression to malignant melanoma with metastases to lymph nodes [61].
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39

40 Genetic characterisation of the UV-induced melanocytic hyperplasias and melanomas
41 of *Monodelphis* found that altered levels of expression of *CDKN2A* and *ARF* genes are
42 associated with the etiology of melanoma formation and progression in these animals [62];
43 both *CDKN2A* and *ARF* are major contributors to melanoma initiation and progression in
44 humans [2]. In addition, *Monodelphis* have both DNA photoactivated excision repair and
45 ubiquitous excision repair mechanisms, as do humans [63]; in contrast, an absence of
46 photoactivation of UV-induced pyrimidine dimers is detectable in mouse epithelial cells in vivo
47 [64]. Studies in *Monodelphis* have also shown that SPF 15 sunscreen is effective in
48 dramatically reducing the incidence of melanocytic naevi [65], and since it was demonstrated
49
50
51
52
53
54
55
56
57
58
59
60

that 43% of *Monodelphis* with nevi progress to malignant melanoma in middle-to-late adulthood [66], this makes *Monodelphis* a model of sunscreen-mediated melanoma prevention.

3.3 Pig

The spontaneous occurrence of melanoma in pigs is generally very low; during a 5-month study of pig carcasses from two abattoirs, only 220/747,014 (0.03%) had gross cutaneous and lymph node lesions suggestive of melanoma [67]. Interestingly, however, 174/176 of the cutaneous lesions submitted for histological analysis were spontaneously regressing melanomas (the remaining 2 were non-regressing melanomas). Through selective breeding to increase the incidence of melanoma in pig breeds, three models of hereditary melanoma in pigs have been established: the Sinclair miniature swine, the Munich miniature swine troll and the Melanoma-bearing Libechov minipig, each showing congenital or early postnatal development of melanoma and spontaneous regression associated with depigmentation.

3.3.1 Sinclair miniature swine

The Sinclair miniature swine was derived from the Hormel miniature pig at the Hormel Institute in the USA. The first observation of cutaneous melanoma in the Sinclair miniature swine strain was in 1967 at an incidence of 21% [68] however with subsequent selective breeding the incidence is now at ~60%, with black coat-colour pigs developing multiple congenitally- or postnatally-appearing skin lesions (exophytic, flat, ulcerated, locally necrotic) and red-coat color developing no tumours [69].

The Sinclair miniature swine melanoma model has numerous features in common with melanoma in humans. Firstly, they develop benign melanocytic lesions (benign naevi) that are capable of malignant transformation (superficial spreading melanoma or nodular melanoma) with metastatic spread of the deeply invasive tumors (to lymph nodes and visceral organs,

mainly lungs and liver), which is analogous to the progression of melanoma seen in humans [70] (Figure 4). Secondly, these melanomas are capable of undergoing spontaneous regression, as has been reported in some human melanoma patients (Figure 4). In addition, the fact that several melanoma lesions can progress and regress simultaneously in the same pig [71] suggests that tumor heterogeneity may play a significant role in the natural history of swine melanoma, similar to that seen in humans. The Sinclair swine melanoma model also affords the unique opportunity to perform sequential biopsies on a single lesion (due to the large size of the swine melanomas) and similarly, the opportunity to perform biopsies on progressing and regressing tumours. Finally, there is a heritable component to the development of melanomas as seen with some melanomas in humans, although as yet the exact genetic determinants responsible are not yet known (structural alterations in chromosomes 2, 3, 6, 7, and 12 may represent the initial step of melanoma development [72]).

Spontaneous regression of melanomas due to a tumour-related immune response has been reported in human patients. In the Sinclair miniature swine, the regression of the melanoma occurs in a two-phase process, with the first phase (typically occurring at 2 months of age) characterised by a massive macrophage infiltration, followed by the second phase (typically occurring at 4 months of age) characterized by lymphocyte infiltration and complete elimination of melanomas [73]. The tumour-infiltrating lymphocytes in the second stage were identified as cytotoxic (CD4-/CD8+), and were found at much higher levels than in the peripheral blood, suggesting that the cytotoxic T-lymphocytes play the main role in the final elimination of the melanoma. However, as antibodies against melanoma antigens were found in the pigs with spontaneously regressing melanoma, usually preceding or appearing together with tumor regression and depigmentation, an antibody-mediated immune response directed against common antigens presented by both malignant and normal melanocytes may also play a role [74]. The spontaneous regression has also been found to be associated with defective telomerase activity [75] and increased sensitivity of the melanoma cells to apoptosis [76].

3.3.2 Munich miniature swine troll

The Munich miniature swine (MMS) troll is historically the second swine model with hereditary melanoma, although it is not as widely used as the other 2 models. It was established at the University of Munich in Germany in 1986, using a melanoma-bearing boar with two unaffected sows from the herd originally developed from the Hanford and the Columbian miniature swine [77]. Similar to the other pig models, selective breeding has led to a high incidence of malignant tumors (~70%), benign melanocytic lesions are also observed, the skin lesions are already present at birth or appear within 2 months after birth, and complete spontaneous regression of melanomas accompanied by hair and skin depigmentation occurs. Interestingly, elevated expression of porcine endogenous retroviruses were detected in melanomas and pulmonary metastasis-derived cell cultures from the MMS troll [78], and upregulated transcription and expression of human endogenous retrovirus (HERV) group HERV-K(HML-2)-encoded proteins has been reported in human melanomas [79]. However, it is not clear what role this retroviral material plays in the onset and/or promotion of tumourigenesis [80].

3.3.3 Melanoma-bearing Libechov minipig

The generation of the melanoma-bearing Libechov minipig model was first started in 1966 at the Czech Academy of Sciences in Libechov by importing Goettingen miniature swine from the University of Goettingen in Germany that were cross-bred with Minnesota miniature pigs from the Hormel Foundation in the USA and Vietnamese pigs from German zoos [81]. Subsequent years of cross-breeding with pigs of several commercial meat breeds and Vietnamese pigs (to maximize genetic variability) produced a population of more than genetically heterogeneous 2,000 descendants, with the first few black piglets with melanoma being observed in 1989. Selective breeding of melanoma-bearing animals confirmed the genetic predisposition to melanoma, with an incidence of ~50% and this new pig model was termed the melanoma-bearing Libechov minipig (MeLiM) [82-83]. Subsequent breeding

1 programs have increased the melanoma incidence to ~80%, with tumour devitalization being
2
3 applied to the very affected pigs to increase their survival and allow their use in breeding [84].
4
5

6 Reflecting the multi-hybrid nature of this strain, MeLiM pigs show variability in coat
7
8 colour, with the black pigs being most affected by melanoma. The black pigs develop three
9
10 types of melanocytic skin lesions (superficial spreading melanoma, nodular melanoma or
11
12 unclassified melanoma) and display a lesion continuum from lentigo to metastatic melanomas
13
14 [85]. The metastases typically occur in the lymph nodes, lungs and spleen (with heavily affected
15
16 animals bearing metastases in other visceral organs). The MeLiM melanomas share a lot of
17
18 similarity with human melanomas, both in terms of their biochemistry and
19
20 immunohistochemical characteristics. For example, the deep-black pigmentation of the
21
22 tumours is due to the high concentration of melanosomes with a high proportion of melanin
23
24 [86], as found in the Sinclair swine [87] and human nodular melanoma [88]. Similarly, increased
25
26 expression of key proteins in melanoma diagnosis and malignancy are found to be elevated in
27
28 both MeLiM melanomas and human melanomas, such as S100, RACK1 and Tenascin C
29
30 (reviewed in [84]). In addition, mass spectrometry analysis of growing MeLiM melanoma tissue
31
32 showed high concentrations of zinc (due to elevated metallothionein content) [89], which has
33
34 also been reported in human melanomas [90].
35
36
37
38
39

40 Similar to the Sinclair miniature swine model, the development of melanoma in the
41
42 MeLiM model is heritable and polygenic. Although the precise genetic determinants not yet
43
44 known, a genome-wide association study (GWAS) revealed several loci on chromosomes 2,
45
46 5, 7, 8, and 16 showing significant associations with melanoma occurrence and progression,
47
48 and comparison to human melanoma GWAS results indicated shared association signals at
49
50 *CDKAL1* and *TERT* loci but also nearby *CCND1*, *FTO*, *PLA2G6* and *TMEM38B-RAD23B* loci
51
52 [91]. Further similarities with the Sinclair miniature swine model include the cutaneous
53
54 melanoma lesions being present either at birth or within first 2 months of life, and the lesions
55
56 either progressing (developing extensive metastasis that leads to death within the first 3
57
58 months of life) or spontaneously regressing (with associated skin and bristle depigmentation)
59
60

[84]. The spontaneous regression occurs asynchronously between the different lesions on the animal (regression is usually completed around 6-12 months of age) and the depigmentation can sometimes spread to the entire body, suggesting activation of immune cells against an antigen common to normal melanocytes as well as melanoma cells [82]. Spontaneous regression in both the Sinclair miniature swine and MeLiM melanomas represent a promising immunological model for monitoring immune cells participating in anti-melanoma reaction. Double-positive (DP) T-lymphocyte ($CD4^+/CD8^{high}$) populations of effector/memory ab T-cells (both in the peripheral blood (PBL) and within the tumour (TIL)) have been reported to expand in MeLiM animals during melanoma regression and it is postulated they are involved in the regression process [92], although this is in contrast to the Sinclair miniature swine model which found the populations of DP lymphocytes were generally consistent in all TIL and PBL populations examined [93]. Nevertheless, a significant increase of DP lymphocytes has also been reported in human melanomas and metastases [94].

Finally, the role of the microbiome in melanoma development has been able to be studied in MeLiM piglets. The number of *Fusobacteria* was higher in melanoma samples compared to healthy skin and also in progressing melanomas compared to regressing ones [95]. Interestingly, the abundance of *Fusobacteria* in the gut is connected with colorectal cancer development and progression in humans (reviewed in [96]), thus it will be of interest to see if the skin microbiome also plays a role in melanoma development or progression in humans.

4. Domesticated animal models of spontaneously occurring melanoma

4.1 Horse

In horses, skin tumours are the most common among neoplasms, with up to 15% of all equine skin tumours being melanocytic [97]. Although >90% of these melanocytic tumours are benign at initial presentation, if left untreated, up to two-thirds can progress to overt malignancy

and are able to undergo widespread metastasis [97-99]. The vast majority appear in grey or white horses, usually at or before the age of 5 years which corresponds to the time in their life when their coat colour changes [97-98; 100-101]. A review of the pathological characteristics of equine melanoma (based on a retrospective study of 53 cases) proposed 4 manifestations of equine melanotic disease: melanocytic nevus, discrete dermal melanoma (benign and malignant forms), dermal melanomatosis, and anaplastic malignant melanoma [102]. The range of intradermal naevi seen in horses resembles those seen in humans, with common melanocytic naevi, cellular blue naevi and combined cellular blue naevi, showing histopathological features in common with their human counterparts [103]. Although discrete dermal melanoma and dermal melanomatosis were described as distinctly separate entities by Valentine [102], these are generally histologically indistinguishable, presenting as indistinct, heavily pigmented tumour cells in the deep dermis, thus it seems equally probable that these diagnoses exist as a continuum [104]. Interestingly, in contrast to melanomas in solid-coloured horses, which although rarer, are characterised by high malignancy and early metastases, melanomas in grey horses tend to have an extended period of benign growth, prior to malignant transformation and metastasis [97; 105]. Nevertheless, grey and white horses are associated with the development of dermal melanomatosis such that beyond the age of 15 years, at least 80% of grey horses will have melanomas at some location [94]. Similar to human melanoma, equine melanomas are typically positive for the S-100, PCNA, HMB-45, Ki-67, T-311 and CD44 markers by immunohistochemistry [101]. In addition, *RACK1* expression, which was originally found to characterise malignant melanocytic lesions in the MeLiM model and, later in human melanomas, has also been found to be effective in distinguishing benign melanocytic tumours from melanomas in horses [106].

The complex inheritance of melanoma and coat/skin pigmentation (grey level, vitiligo grade, and speckling grade) in grey horses **has** been shown to be primarily due to the effects of a 4.6-kb duplication in intron 6 of the *STX17* (*syntaxin 17*) gene; this constitutes a cis-acting regulatory mutation that has melanocyte-specific effects (possibly due to the fact that it affects

an enhancer that encodes binding sites for the microphthalmia-associated transcription factor (*MITF*), which regulates melanocyte development) [107-108]. Both *STX17* and the neighbouring *NR4A3* (*nuclear receptor subfamily 4, group A, member 3*) gene are overexpressed in melanomas from grey horses. Elevated *STX17* is associated with constitutive activation of the ERK pathway in melanocytic cells [109], thus highlighting the similarities to human melanoma in which the MAPK/ERK pathway is also involved in melanomagenesis. Similarly, just as MC1R plays an important role in regulating skin pigmentation and melanoma growth in humans, and expression of agouti signalling protein (*ASIP*), known MC1R antagonist, slows B16-F10 melanoma growth and increases survival times in mice [110], grey horses carrying a loss-of-function mutation in *ASIP* have a higher incidence of melanoma [107], suggesting that increased melanocortin-1 receptor signalling promotes melanoma development in grey horses, as it does in humans.

Melanoma nodules occur most frequently underneath the tail and at high rates in the peri-anal region, lips, and eyelids, with some also noted in the vulva [105; 111] (Figure 5). As many of these areas are near mucosal epithelium, equine melanomas may represent a model for mucosal melanoma in humans. Next-generation sequencing of equine melanomas that originated from mucosal-like (perineum, perianal region, prepuce, vulva, or ventral tail) or mucocutaneous sites (near the eyes or mouth, and show both mucosa and haired skin) were whole-exome sequenced, as well as some from cutaneous sites (haired skin only) and other sites (urinary bladder wall muscle and the parotid gland), found that similar to human mucosal melanomas, most equine melanomas had less than 5 mutations/Mb [112], unlike UV-associated human subtypes such as superficial spreading and nodular melanoma [113]. In addition, the most prominent driver genes were *NRAS* and *TP53*, as has been reported in human mucosal melanoma samples [112]. Both species also showed mutations in *PTEN* and *KIT*, as well as a few cases with *BRAF* mutations. Interestingly however, equine melanoma cases from mucosal-like or mucocutaneous sites, showed a landscape of driver genes that was less populated than that seen for human mucosal melanoma, and there were fewer

1 recurrently mutated genes in common with human mucosal melanoma, with no mutations
2 found in *SF3B1*, *ATRX* or *NF1* [112]. The equine cutaneous melanomas had no point mutations
3 or indels in known human melanoma driver genes (apart from one having an *NRAS*^{Q61R}
4 mutation) [112], and as such they show some similarities to pigmented epithelioid
5 melanocytoma ('animal-type melanoma'), a rare type of melanoma in humans that does not
6 carry mutations in genes frequently mutated in cutaneous melanoma [114].
7
8
9
10
11
12
13
14
15
16

17 **4.2 Cat**
18
19
20
21

22 Melanoma of any site in the cat is rare, with only 4 melanomas found out of 3,145 feline
23 diagnoses at the Animal Medical Center in New York [115]. However, in contrast to other
24 domesticated species such as dogs and horses, cats develop intraocular melanoma more
25 frequently than cutaneous melanoma [116]. In a study of 29 feline melanomas over an 11 year
26 period, 19 melanomas were intraocular, with the iris being the most common area of
27 involvement, metastasis occurring in 63% of these cases, and 62.5% of the cats requiring
28 euthanasia ~5 months post-diagnosis [116]. Indeed, iris melanoma is the most frequent feline
29 intraocular tumour (Figure 6), with only rare cases of choroidal melanoma being reported [117].
30 This is in contrast to humans, where most intraocular (uveal) melanomas originate in the
31 choroid (with iris melanoma being rare); these do not usually spread and are associated with
32 a >95% 5-year relative survival rate. Nevertheless, these feline ocular melanomas may offer
33 an alternative to the other animal models of ocular melanoma, which suffer the limitations that
34 they do not replicate the critical behaviour of the disease in humans, specifically spontaneous
35 occurrence and concurrent metastasis (reviewed in [118]).
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55

56 **4.3 Dog**
57
58
59
60

Melanocytic tumours are relatively common in the dog, and are typically malignant; a study of 2,350 cases of melanocytic tumours found 70% were malignant melanomas and 30% were benign tumours (melanocytomas) [119]. The anatomical locations of melanoma in the dog are oral (62% of cases), cutaneous (27%), digit (6%), ungual (4%) and ocular (1%) [119]. Interestingly, the correlation with malignancy varies at each site, with 84-100% of oral, digit and ungual melanocytic tumours being malignant (melanoma), in contrast to only 43% and 29% of cutaneous and ocular melanocytic tumours being malignant (respectively) [119].

4.3.1 Mucosal melanoma

Canine melanomas of the oral cavity occur mostly on the gingiva, but also on the lips, tongue, tonsils, palate, and oropharynx (Figure 7a) [119-120]. Spontaneous anal sac, intestinal and intranasal melanoma have also been reported, but these mucosal sites are much more rare [121]. Similar to humans, epidemiological studies have shown certain breeds more prone (at a higher-risk) to develop oral melanoma, with Poodles, Golden Retrievers, Labradors, Rottweilers and Yorkshire terriers representing 50% of the cases [121]. Similar to human mucosal melanoma, canine oral melanoma is an aggressive tumour showing local invasiveness and a high propensity to metastasize (to regional lymph nodes and lungs) [122], with reported median survival times of 3-24 months (depending on stage and treatment) [119]. Further parallels with their human counterpart is the fact that canine melanomas are resistant to chemotherapy and radiation therapy [123]. There are also similarities between the species in terms of response to therapy. Clinical trials showed that patients with metastatic mucosal melanoma can benefit from treatment with the immunotherapy regimes of Nivolumab (targeted PD-1) alone or in combination with Ipilimumab (targeting CTLA-4) [124] and similarly, a rat-dog chimaeric anti-PD-L1 monoclonal antibody was well tolerated and showed some clinical efficacy in the treatment of canine oral malignant melanoma with a response rate of 14.3% [125]. In addition, the use of human genes has also proved beneficial in treating canine oral

1 melanoma. For example, in a phase II clinical trial using allogeneic whole-cell vaccination, a
2 canine melanoma cell line transfected with human *gp100*, killed by irradiation and intradermally
3 administered to dogs with malignant melanoma, found objective evidence of tumour regression
4 in 17.6% of cases, with dogs experiencing tumour control surviving significantly longer (337
5 days) than dogs having no response (95 days) [126]. Similarly, intratumoral administration of
6 the human *CD94L* gene was evaluated in a phase I clinical trial in four dogs with oral melanoma
7 and in three of the dogs, a 12-58% reduction of tumour burden was reported, with no adverse
8 effects observed [127]. Finally, a bacterial plasmid DNA vaccine encoding the human
9 tyrosinase antigen (Oncept™) is licensed for the adjuvant treatment of stage II and III canine
10 oral melanoma after loco-regional control and is currently awaiting USDA approval for use in
11 horses with melanoma. However, it should be noted that there is controversy about its efficacy
12 as some studies have observed responses in dogs with macroscopic disease and suggested
13 the vaccine could be considered as palliative treatment in dogs with stage IV disease [128],
14 whilst others have found that dogs who received the vaccine did not achieve a greater
15 progression-free survival, disease-free interval or median survival time than dogs that did not
16 receive the vaccine [129].

17
18 A study that compared the histopathological features of human mucosal melanoma and
19 canine oral melanoma found analogous architectural features were important for diagnosis and
20 staging of the melanoma in both species [130]. Interestingly, both human and canine mucosal
21 melanomas included the range of epithelioid, spindloid, mixed epithelioid/spindloid, or small
22 round blue cell melanocyte morphologies [130]. Also noteworthy was the frequent presence of
23 a lentiginous intra-epithelial component in the canine melanomas, which is a feature known to
24 precede invasive melanomas in humans that subsequently metastasize, thus indicating a
25 similar pathway of progression in the dog [130].

26
27 At the molecular level, there are many similarities of canine oral melanoma with human
28 mucosal melanoma. Next-generation sequencing of canine oral melanomas revealed they had
29 less than 5 mutations/Mb, similar to human mucosal and equine mucosal melanomas [112]. In

1 addition, the most prominent driver genes were *NRAS* and *TP53*, as has been reported in
2 human mucosal melanoma samples, with both species also showing mutations in *NF1*, *PTEN*,
3 *KRAS* and *BRCA2* [112; 131]. **Importantly**, both species show involvement of the MAPK and
4 PI3K/AKT pathways in mucosal melanoma formation and progression [112; 131-132], with
5 constitutive pathway activation and similar sensitivity to AZD6244 and rapamycin observed in
6 both in human melanoma and canine oral melanoma cell lines [133], suggesting that canine
7 oral melanoma could be a good model for human mucosal melanoma. Although they did not
8 show any mutations in other drivers of mucosal melanoma frequently in human patients, such
9 as *SF3B1* and *ATRX*, suggesting that these canine cancers may not represent a faithful model
10 for the subset of human mucosal melanoma patients with mutations in these genes. Yet, some
11 canine oral melanomas showed mutations in *EIF1AX* (G9V and R13C), an essential translation
12 initiation factor that is mutated in human uveal melanoma [8] and meningeal melanocytic
13 tumours [134], with the same N-terminal amino acid substitutions at conserved sites G9 and
14 R13 also having been found in multiple other human tumours (COSMIC database:
15 COSM6908971, COSM5899335) [112]; thus the canine oral melanomas have helped to
16 identify the importance of these conserved regions in human tumourigenesis. In addition,
17 cross-species comparative copy number analysis found that the most significant deletion in
18 human mucosal melanoma samples was a deletion at 15q15.1, which is a region syntenic to a
19 significant deletion in canine oral melanomas found on canine chromosome 30; both species
20 harboured *BUB1B*, *KNSTRN* and *B2M* within this deletion regions and these genes are
21 frequently altered in human tumours [112]. Similarly, both *MDM2* and *SMO* were found to be
22 amplified in human mucosal melanomas and canine oral melanomas, with these genes having
23 known roles in cutaneous melanoma and a range of cancers, respectively [112]. In addition, a
24 recent study looking at long non-coding RNAs (lncRNAs) in canine oral melanomas found one
25 down-regulated (*SOX21-AS*) and two up-regulated (*CASC15* and *ZEB2-AS*) differentially
26 expressed lncRNAs ("onco-lncRNAs"), that have been shown to be differentially expressed in
27 a range of human cancers [135].

1
2
3
4 4.3.2 Cutaneous melanoma

5
6 Cutaneous melanocytic tumours in dogs are usually dermal in origin and are typically
7
8 benign, referred to as melanocytomas [136]. The most common sites for benign cutaneous
9
10 melanomas are the face (near the eyelids), trunk, and extremities, and malignant melanoma is
11
12 found most frequently on the head, ventral abdomen, and scrotum, with metastases primarily
13
14 occurring to the lymph nodes and lungs [136]. Cutaneous melanoma occurs more commonly
15
16 in dogs with heavily pigmented skin, with Schnauzers and Scottish terriers at increased risk
17
18 [137], which is in contrast to cutaneous melanoma in humans where individuals of light/fair skin
19
20 are at increased risk.
21
22

23
24 There is limited data available on the somatic alterations of canine cutaneous
25
26 melanoma. One study took a candidate gene approach with 20 cutaneous melanomas and
27
28 found no variants mutations in the key melanoma drivers, specifically *BRAF*, *NRAS*, *PTEN*,
29
30 *KIT*, *GNAQ*, and *CDK4* [119], whereas another study on 2 dogs found one carried an *NRAS*
31
32 mutation and one carried *KRAS*, *TP53* and *KIT* mutations [138]. These reports, taken together
33
34 with the fact that most canine cutaneous melanocytic tumours are benign and occur on hair-
35
36 bearing skin, thus are not linked to UV-light exposure, suggest these canine cutaneous
37
38 melanomas arise from a distinct mechanism, different from that of most human cutaneous
39
40 melanomas, and as such may be of more relevance to acral melanomas or other rare
41
42 melanomas, and as such may be of more relevance to acral melanomas or other rare
43
44 categories of dermal melanomas in humans [2].
45
46
47
48

49 4.3.3 Ocular melanoma

50
51
52
53
54 Melanocytic tumours are the most common ocular neoplasms in dogs, and arise most
55
56 frequently in the anterior uvea; a study of intraocular melanocytic tumours in 244 dogs reported
57
58 the anterior uvea as the tumour site in 85% of cases, with the limbic area and choroid
59
60 accounting for 13 and 5%, respectively [139]. Similar to cutaneous melanocytic tumours, most

1 intraocular melanocytic tumours in dogs are benign ('benign uveal melanoma'; Figure 7b), with
2 reports of malignancy ranging from 15-30%, and those with metastatic behaviours accounting
3 for ~4% [136]. The prognosis for dogs with histologically benign uveal melanoma is excellent,
4 however, those with malignant uveal melanoma have significantly shorter lifespans [139].
5 Interestingly, expression profiling using a 12-gene assay has been shown to reliably distinguish
6 metastasising from non-metastasising uveal melanomas in humans [140], and four of these
7 genes (*HTR2B*, *FXR1*, *LTA4H*, and *CDH1*) have also been found to be overexpressed in
8 metastasising canine uveal melanoma [141].
9
10
11
12
13
14
15
16
17
18
19
20
21

22 4.3.4 Acral melanoma

23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
Melanoma is the second most common digital tumour in dogs, accounting for 15-17%
of neoplasias of the digits [142-143]. Similar to human acral melanoma, which arises on the
plantar surface of the foot, palms of hands, and fingers, canine acral melanoma arises on the
footpad or nail bed. Studies have shown that most canine melanocytic tumors of the digits are
malignant (49-86% of cases) and all nail bed melanocytic tumors are malignant [119]. As with
other forms of melanoma in the dog, the development of melanoma in the digits of dogs are
over-represented in some breeds, such as the Scottish terrier, Schnauzer, Beauce shepherds,
and Rottweilers [142-143]. These tumours are locally aggressive, with bone lysis occurring in
40-58% of cases and a high propensity to metastasize, with lymph node or lung metastasis
found in 30-40% of cases at the time of presentation [142]. To date, only 3 melanomas from
the digits of dogs have been characterized at the genomic level with 2 having *KRAS* mutations
and 1 having an *NRAS* mutation [131].

56 4.3.5 Leptomeningeal melanoma

1 There has been little research into leptomeningeal melanoma/melanomatosis in dogs,
2
3 most probably due to the rarity of the disease. Indeed, there is only one published report of
4
5 primary CNS melanoma in a dog; an 11-year old Black and Tan Coonhound with disseminated
6
7 melanoma involving the right femoral bone marrow, lung, multiple lymph nodes, and adrenal
8
9 gland, with diffuse infiltration of the leptomeninges of the brain and spinal cord, and in light of
10
11 the lesion distribution that resembled leptomeningeal melanomatosis in humans, the diagnosis
12
13 was given as primary leptomeningeal melanomatosis with multiorgan metastasis [144]. Similar
14
15 to humans, secondary (metastatic) CNS melanoma is more common than primary CNS
16
17 melanoma; post-mortem studies of dogs with secondary intracranial tumours found them to be
18
19 malignant melanomas in ~3.4% of cases [145-146].
20
21
22
23
24
25

26 **5. Conclusion**
27
28
29
30

31 The key feature(s) of each animal as a 'model' of melanoma in humans is listed in Table
32
33 3. No one animal is the perfect model, however, the spontaneous nature of the development
34
35 of melanoma and its metastatic spread in the platyfish/swordtail, opossum, miniature pig,
36
37 horse, cat and dog, as well as the genetic heterogeneity of the latter three species, and the
38
39 additional advantage of a shared microenvironmental exposures in companion animals, surely
40
41 makes these animals attractive candidate models.
42
43
44
45
46
47
48

49 **Acknowledgments**
50
51
52
53

54 DJA is supported by Cancer Research UK (21717), the Medical Research Council
55
56 (MR/S01473X/1) and the Wellcome Trust. EEP is funded by the MRC HGU Programme
57
58 (MC_UU_00007/9), European Research Council (ZF-MEL-CHEMBIO-648489), and L'Oreal-
59
60 Melanoma Research Alliance (401181).

Author contribution statement

LvdW and TB wrote the section of the manuscript on the genetics and pathology of human melanoma. LvdW, EEP and GAW wrote the sections of the manuscript on the occurrence of melanoma in animals. EEP, GAW and DJA co-ordinated obtaining the photographs. All authors contributed to revision of the manuscript and the final published paper. All authors contributed equally.

For Peer Review

Reference list

1. Chapter 2: Melanocytic tumors. In *WHO Classification of Skin Tumors*, (4th edn), Elder DE, Massi D, Willemze R, Scolyer R (eds). International Agency for Research on Cancer: Lyon, 2018; 470 pages.
2. Bastian BC. The molecular pathology of melanoma: an integrated taxonomy of melanocytic neoplasia. *Annu Rev Pathol* 2014; **9**: 239–271.
3. Linos E, Swetter SM, Cockburn MG, *et al*. Increasing burden of melanoma in the United States. *J Investig Dermatol* 2009; **129**: 1666–1674.
4. Larkin J, Chiarion-Sileni V, Gonzalez R, *et al*. Five-year survival with combined nivolumab and ipilimumab in advanced melanoma. *NEJM* 2019; **381**:1535–1546.
5. Yde SS, Sjoegren P, Heje M, *et al*. Mucosal Melanoma: a Literature Review. *Curr Oncol Rep* 2018; **20**: 28.
6. Mihajlovic M, Vlajkovic S, Jovanovic P, *et al*. Primary mucosal melanomas: a comprehensive review. *Int J Clin Exp Pathol* 2012; **5**: 739–753.
7. Williams MD. Update from the 4th Edition of the World Health Organization Classification of Head and Neck Tumours: Mucosal Melanomas. *Head Neck Pathol* 2017; **11**: 110–117.
8. Robertson AG, Shih J, Yau C, *et al*. Integrative Analysis Identifies Four Molecular and Clinical Subsets in Uveal Melanoma. *Cancer Cell* 2017; **32**:204–220.
9. Shields CL, Markowitz JS, Belinsky I, *et al*. Conjunctival melanoma: outcomes based on tumor origin in 382 consecutive cases. *Ophthalmology* 2011; **118**: 389–395.
10. Kenawy N, Lake SL, Coupland SE, *et al*. Conjunctival melanoma and melanocytic intra-epithelial neoplasia. *Eye (Lond)* 2013; **27**: 142–152.
11. Küsters-Vandeveld HV, Küsters B, van Engen-van Grunsven AC, *et al*. Primary melanocytic tumors of the central nervous system: a review with focus on molecular aspects. *Brain Pathol* 2015; **25**:209–226.

12. Helseth A, Helseth E, Unsgaard G. Primary meningeal melanoma. *Acta Oncol* 1989; **28**:103–104.
13. Ferlay J, Shin H-R, Bray F, *et al.* Estimates of worldwide burden of cancer in 2008: GLOBOCAN 2008. *Int J Cancer* 2010; **127**: 2893–2917.
14. Amer MH, Al-Sarraf M, Baker LH, *et al.* Malignant melanoma and central nervous system metastases: incidence, diagnosis, treatment and survival. *Cancer* 1978; **42**: 660–668.
15. Jellinger K, Böck F, Brenner H. Meningeal melanocytoma. Report of a case and review of the literature. *Acta Neurochir (Wien)* 1988; **94**: 78–87.
16. Helseth A, Helseth E, Unsgaard G. Primary meningeal melanoma. *Acta Oncol* 1989; **28**: 103–104.
17. Kadonaga JN, Frieden IJ. Neurocutaneous melanosis: definition and review of the literature. *J Am Acad Dermatol* 1991; **24**:747–755.
18. Mitre V, Heym K, Clark GD, *et al.* Neurocutaneous Melanocytosis and Leptomeningeal Melanoma. *J Pediatr Hematol Oncol* 2019; DOI:10.1097/MPH.0000000000001680
19. Van Beek EJA, Balm AJM, Nieweg OE, *et al.* Treatment of regional metastatic melanoma of unknown primary origin. *Cancers (Basel)* 2015; **7**:1543–1553.
20. Pfeil AF, Leiter U, Buettner PG, *et al.* Melanoma of unknown primary is correctly classified by the AJCC melanoma classification from 2009. *Melanoma Res* 2011; **21**: 228–234.
21. Tos T, Klyver H, Drzewiecki KT. Extensive screening for primary tumor is redundant in melanoma of unknown primary. *J Surg Oncol* 2011; **104**: 724–727.
22. Blessing K, McLaren KM. Histological regression in primary cutaneous melanoma: recognition, prevalence and significance. *Histopathology* 1992; **20**: 315–322.
23. Egberts F, Bergner I, Kruger S, *et al.* Metastatic melanoma of unknown primary resembles the genotype of cutaneous melanomas. *Ann Oncol* 2014; **25**: 246–250.
24. Perez Guijarro E, Day CP, Merlino G., *et al.* Genetically Engineered Mouse Models of Melanoma. *Cancer* 2017; **123**: 2089–2103

25. Richards JR, Yoo JH, Shin D, *et al.* Mouse models of uveal melanoma: Strengths, weaknesses, and future directions. *Pigment Cell Melanoma Res* 2020; **33**: 264–278.
26. van der Weyden L, Arends MJ, Brenn T, *et al.* Widespread spontaneous hyperproliferation, melanosis and melanoma in Hgf-Cdk4R24C mice. *Melanoma Res* 2018; **28**: 76–78.
27. Moore AR, Ran L, Guan Y, *et al.* GNA11 Q209L Mouse Model Reveals RasGRP3 as an Essential Signaling Node in Uveal Melanoma. *Cell Rep* 2018; **22**: 2455–2468.
28. Walker GJ, Hayward NK. Pathways to melanoma development: lessons from the mouse. *J Invest Dermatol* 2002; **119**: 783–792.
29. van der Weyden L, Patton EE, Wood GA, *et al.* Cross-species models of human melanoma. *J Pathol* 2016; **238**: 152–165.
30. Larue, L. Origin of Mouse Melanomas. *J Invest Dermatol* 2012; **132**: 2135–2136.
31. Waldoch JA, Burke SS, Ramer JC *et al.* Melanoma in the Skin of a Nurse Shark (*Ginglymostoma Cirratum*). *J Zoo Wildl Med* 2010; **41**: 729–731.
32. Nau MR, Gardiner DW, Nilson E, *et al.* Cutaneous malignant melanoma in a Haller's round ray *Urobatis halleri*. *Dis Aquat Organ* 2016; **120**: 245–250.
33. Sweet M, Kirkham N, Bendall M, *et al.* Evidence of melanoma in wild marine fish populations. *PLoS One* 2012; **7**: e41989.
34. Elkan E. Malignant melanoma in a snake. *J Comp Pathol* 1974; **84**: 51–57.
35. Reid HAC, Herron AJ, Hines II ME, *et al.* Metastatic Malignant Melanoma in a Mandarin Duck (*Aix galericulata*). *Avian Diseases* 1993; **37**: 1158–1162.
36. Kufuor-Mensah E, Watson GL. Malignant melanomas in a penguin (*Eudyptes chrysolophus*) and a red-tailed hawk (*Buteo jamaicensis*). *Vet Pathol* 1992; **29**: 354–356.
37. Ueda K, Ueda A, Ozaki K. Cutaneous malignant melanoma in two rabbits (*Oryctolagus cuniculus*). *J Vet Med Sci* 2018; **80**: 973–976.

38. Mavangira V, Hughes JM, Middleton JR, et al. Malignant melanoma of the horn base in a Pygora goat. *J Vet Diagn Invest* 2008; **20**:104–107.
39. Mollat WH, Gailbreath KL, Orbell GM. Metastatic malignant melanoma in an alpaca (*Vicugna pacos*). *J Vet Diagn Invest* 2009; **21**: 141–144.
40. Saunders RA, Killick RS, Barrows MG, et al. Diagnosis and treatment of concurrent dermal malignant melanoma and melanocytomas in a pygmy hippopotamus (*Choeropsis liberiensis*). *Vet Dermatol* 2017; **28**: 520–e127.
41. Steeil JC, Schumacher J, Baine K, et al. Diagnosis and treatment of a dermal malignant melanoma in an African lion (*Panthera leo*). *J Zoo Wildl Med* 2013; **44**: 721–727.
42. Winslow CM, Wood J, Gilliam JN, et al. Congenital amelanotic melanoma in a crossbred heifer calf. *J Vet Diagn Invest* 2017; **29**: 544–547.
43. Ameri M, Anderson WI, Orlando HM. Cutaneous melanocytoma in a cynomolgus monkey (*Macaca fascicularis*): cytologic and histologic characteristics. *Comparative Clinical Pathology* 2012; **21**: 202–207.
44. Heckers KO, Aupperle H, Schmidt V, et al. Melanophoromas and iridophoromas in reptiles. *J Comp Pathol* 2012; **146**: 258–268.
45. Meierjohann S, Scharl M. From Mendelian to molecular genetics: the Xiphophorus melanoma model. *Trends Genet* 2006; **22**: 654–661.
46. Wittbrodt J, Adam D, Malitschek B et al. Novel Putative Receptor Tyrosine Kinase Encoded by the Melanoma-Inducing Tu Locus in Xiphophorus. *Nature* 1989; **341**: 415–421.
47. Regneri J, Klotz B, Wilde B, et al. Analysis of the putative tumor suppressor gene *cdkn2ab* in pigment cells and melanoma of Xiphophorus and medaka. *Pigment Cell Melanoma Res* 2019; **32**: 248–258.
48. Fernandez AA, Morris MR. Mate choice for more melanin as a mechanism to maintain a functional oncogene. *Proc Natl Acad Sci U S A* 2008; **105**: 13503–13507.

49. Fernandez AA. A cancer-causing gene is positively correlated with male aggression in *Xiphophorus cortezi*. *J Evol Biol* 2010; **23**: 386–396.
50. Fernandez AA, Bowser PR. Selection for a dominant oncogene and large male size as a risk factor for melanoma in the *Xiphophorus* animal model. *Mol Ecol* 2010; **19**: 3114–3123.
51. Kazianis S, Morizot DC, Coletta LD, *et al.* Comparative structure and characterization of a CDKN2 gene in a *Xiphophorus* fish melanoma model. *Oncogene* 1999; **18**: 5088–5099.
52. Solus JF, Kraft S. Ras, Raf, and MAP kinase in melanoma. *Adv Anat Pathol* 2013; **20**: 217–226.
53. Chin L, Pomerantz J, Polsky D, *et al.* Cooperative effects of INK4a and ras in melanoma susceptibility in vivo. *Genes Dev* 1997; **11**: 2822–2834.
54. Morcinek JC, Weisser C, Geissinger E, *et al.* Activation of STAT5 triggers proliferation and contributes to anti-apoptotic signalling mediated by the oncogenic Xmrk kinase. *Oncogene* 2002; **21**: 1668–1678.
55. Hassel JC, Winnemöller D, Scharl M, *et al.* STAT5 contributes to antiapoptosis in melanoma. *Melanoma Res.* 2008; **18**: 378–385.
56. Setlow RB, Woodhead AD, Grist E. Animal Model for Ultraviolet Radiation-Induced Melanoma: Platyfish-Swordtail Hybrid. *Proc Natl Acad Sci USA* 1989; **86**: 8922–8926.
57. Lu Y, Boswell M, Boswell W, *et al.* Comparison of *Xiphophorus* and human melanoma transcriptomes reveals conserved pathway interactions. *Pigment Cell Melanoma Res* 2018; **31**: 496–508.
58. VandeBerg JL, Robinson ES. The laboratory opossum (*Monodelphis domestica*) in laboratory research. *ILAR J* 1997; **38**: 4–12.
59. Ley RD, Applegate LA, Padilla RS, *et al.* Ultraviolet Radiation-Induced Malignant Melanoma. *Photochem Photobiol* 1989; **50**: 1–5.

60. Kusewitt DF, Applegate LA, Ley RD. Ultraviolet radiation-induced skin tumors in a South American opossum (*Monodelphis domestica*). *Vet Path* 1991; **28**: 55–65.
61. Robinson ES, VandeBerg JL, Hubbard GB, *et al*. Malignant melanoma in ultraviolet irradiated laboratory opossums: Initiation in suckling young, metastasis in adults, and xenograft behavior in nude mice. *Cancer Res* 1994; **54**: 5986–5991.
62. Chan J, Robinson ES, Atencio J, *et al*. Characterization of the CDKN2A and ARF genes in UV-induced melanocytic hyperplasias and melanomas of an opossum (*Monodelphis domestica*). *Mol Carcinog* 2001; **31**: 16–26.
63. Ley RD, Applegate LA. Ultraviolet radiation-induced histopathologic changes in the skin of the marsupial *Monodelphis domestica*. II. Quantitative studies of the photoreactivation of induced hyperplasia and sunburn cell formation. *J Invest Dermatol* 1985; **85**: 365–367.
64. Ley RD, Sedita BA, Grube DD. Absence of photoreactivation of pyrimidine dimers in the epidermis of hairless mice following exposures to ultraviolet light. *Photochem Photobiol* 1978; **27**: 483–485.
65. Nair HB, Ford A, Dick EJ Jr, *et al*. Modeling sunscreen-mediated melanoma prevention in the laboratory opossum (*Monodelphis domestica*). *Pigment Cell Melanoma Res* 2014; **27**: 843–845.
66. Robinson ES, Hubbard GB, Colon G, *et al*. Low-dose ultraviolet exposure early in development can lead to widespread melanoma in the opossum model. *Int J Exp Pathol* 1998; **79**: 235–244.
67. Bundza A, Feltmate TE. Melanocytic cutaneous lesions and melanotic regional lymph nodes in slaughter swine. *Can J Vet Res* 1990; **54**: 301–304.
68. Strafuss AC, Dommert AR, Tumbleson ME, *et al*. Cutaneous Melanoma in Miniature Swine. *Lab Anim Care* 1968; **18**: 165–169.
69. Hook RR, Aultman MD, Adelstein EH, *et al*. Influence of Selective Breeding on the Incidence of Melanomas in Sinclair Miniature Swine. *Int J Cancer* 1979; **24**: 668–672.

- 1 70. Oxenhandler RW, Adelstein EH, Haigh JP, *et al.* Malignant melanoma in the Sinclair
2 miniature swine: An autopsy study of 60 cases. *Am J Pathol* 1979; **96**: 707–720.
3
4
5 71. Hook RR Jr, Aultman MD, Millikan LE, *et al.* The biology of cutaneous exophytic
6 melanomas in Sinclair (S-1) miniature swine. *Am Assoc Cancer Res* 1977; **18**: 46.
7
8
9 72. Pathak S, Amoss MS. Genetic Predisposition and Specific Chromosomal Defects
10 Associated with Sinclair Swine Malignant Melanomas. *Int J Oncol* 1997; **11**: 53–57.
11
12
13 73. Greene JF, Townsend JS, Amoss MS. Histopathology of Regression in Sinclair Swine
14 Model of Melanoma. *Lab. Investig J Tech Methods Pathol* 1994; **71**: 17–24.
15
16
17 74. Cui J, Chen D, Misfeldt ML, *et al.* Antimelanoma Antibodies in Swine with
18 Spontaneously Regressing Melanoma. *Pigment Cell Res* 1995; **8**: 60–63.
19
20
21 75. Pathak S, Multani AS, McConkey DJ, *et al.* Spontaneous Regression of Cutaneous
22 Melanoma in Sinclair Swine is Associated with Defective Telomerase Activity and
23 Extensive Telomere Erosion. *Int J Oncol* 2000; **17**, 1219–1243.
24
25
26 76. Gossett R, Kier AB, Schroeder F, *et al.* Cycloheximide-Induced Apoptosis in Melanoma
27 Cells Derived from Regressing Cutaneous Tumours of SINCLAIR Swine. *J Comp Pathol*
28 1996; **115**: 353–372.
29
30
31 77. Muller S, Wanke R, Distl O. Inheritance of Melanocytic Lesions and Their Association
32 with the White Colour Phenotype in Miniature Swine. *J Anim Breed Genet* 2001; **118**:
33 275–283.
34
35
36 78. Dieckhoff B, Puhlmann J, Büscher K, *et al.* Expression of porcine endogenous
37 retroviruses (PERVs) in melanomas of Munich miniature swine (MMS) Troll. *Vet*
38 *Microbiol* 2007; **123**: 53–68.
39
40
41 79. Schmitt K, Reichrath J, Roesch A, *et al.* Transcriptional profiling of human endogenous
42 retrovirus group HERV-K(HML-2) loci in melanoma. *Genome Biol Evol* 2013; **5**: 307–
43 328.
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
80. Downey RF, Sullivan FJ, Wang-Johanning F, *et al.* Human endogenous retrovirus K and cancer: Innocent bystander or tumorigenic accomplice? *Int J Cancer* 2015; **137**: 1249–1257.
81. Glodek, P. Breeding Program and Population Standards of the Goettingen Miniature Swine. In *Swine in Biomedical Research (Vol 1)*, Tumbleson ME (ed). Plenum Press: New York, 1986; 23–28.
82. Horak V, Fortyn K, Hruban V, *et al.* Hereditary Melanoblastoma in Miniature Pigs and its Successful Therapy by Devitalization Technique. *Cell Mol Biol Noisy Gd Fr* 1999; **45**: 1119–1129.
83. Hruban V, Horak V, Fortyn K, *et al.* Inheritance of Malignant Melanoma in the MeLiM Strain of Miniature Pigs. *Vet Med (Praha)* 2004; **49**: 453–459.
84. Horak V, Palanova A, Cizkova J, *et al.* Melanoma-Bearing Libechov Minipig (MeLiM): The Unique Swine Model of Hereditary Metastatic Melanoma. *Genes (Basel)* 2019; **10**: E915.
85. Vincent-Naulleau S, Le Chalony C, Leplat JJ, *et al.* Clinical and histopathological characterization of cutaneous melanomas in the melanoblastoma-bearing Libechov minipig model. *Pigment Cell Res* 2004; **17**:24–35.
86. Borovansky J, Horak V, Elleder M, *et al.* Biochemical Characterization of a New Melanoma Model-The Minipig MeLiM Strain. *Melanoma Res* 2003; **13**: 543–548.
87. Millikan LE, Hook RR, Manning PJ. Immunobiology of Melanoma. Gross and Ultrastructural Studies in a New Melanoma Model: The Sinclair Swine. *Yale J Biol Med* 1973; **46**: 631–645.
88. Hunter JA, Zaynoun S, Paterson WD, *et al.* Cellular Fine Structure in the Invasive Nodules of Different Histogenetic Types of Malignant Melanoma. *Br J Dermatol* 1978; **98**: 255–272.

89. Anyz J, Vyslouzilova L, Vaculovic T, *et al.* Spatial Mapping of Metals in Tissue-Sections Using Combination of Mass-Spectrometry and Histology Through Image Registration. *Sci Rep* 2017; **7**: 40169.
90. Gorodetsky R, Sheski J, Weinreb A. Iron, Copper, and Zinc Concentrations in Normal Skin and in Various Nonmalignant and Malignant Lesions. *Int J Dermatol* 1986; **25**: 440–445.
91. Bourneuf E, Estelle J, Blin A, *et al.* New Susceptibility Loci for Cutaneous Melanoma Risk and Progression Revealed Using a Porcine Model. *Oncotarget* 2018; **9**: 27682–27697.
92. Cizkova J, Sinkorova Z, Strnadova K, *et al.* The Role of $\alpha\beta$ T-Cells in Spontaneous Regression of Melanoma Tumors in Swine. *Dev. Comp. Immunol.* 2019, **92**, 60–68.
93. Morgan CD, Measel JW Jr, Amoss MS Jr, *et al.* Immunophenotypic characterization of tumor infiltrating lymphocytes and peripheral blood lymphocytes isolated from melanomatous and non-melanomatous Sinclair miniature swine. *Vet Immunol Immunopathol* 1996; **55**:189–203.
94. Desfrancois J, Moreau Aubry A, Vignard V, *et al.* Double Positive CD4CD8 Alphabeta T Cells: A New Tumor-Reactive Population in Human Melanomas. *PLoS ONE* 2010; **5**: e8437.
95. Mrazek J, Mekadim C, Kucerovala P, *et al.* Melanoma-Related Changes in Skin Microbiome. *Folia Microbiol (Praha)* 2019; **64**: 435–442.
96. Zhou Z, Chen J, Yao H, *et al.* Fusobacterium and Colorectal Cancer. *Front Oncol* 2018; **8**: 371.
97. Johnson PJ. Dermatologic tumors (excluding sarcoids). *Vet Clin North Am Equine Pract* 1998; **14**: 625–658.
98. Levene A. Equine melanotic disease. *Tumori* 1971; **57**: 133–168.
99. Schott HC, Major MD, Grant BD, *et al.* Melanoma as a cause of spinal cord compression in two horses. *J Am Vet Med Assoc* 1990; **196**: 1820–1822.

100. McFadyean S: Equine melanomatosis. *J Comp Med* 1933; **46**: 186–204.
101. Seltenhammer MH, Heere-Ress E, Brandt S, *et al.* Comparative histopathology of grey-horse-melanoma and human malignant melanoma. *Pigment Cell Res* 2004; **17**: 674–681.
102. Valentine BA. Equine Melanocytic Tumors: A Retrospective Study of 53 Horses (1988 to 1991). *J Vet Intern Med* 1995; **9**: 291–297.
103. Schöniger S, Summers BA. Equine skin tumours in 20 horses resembling three variants of human melanocytic naevi. *Vet Dermatol* 2009; **20**:165–173.
104. Moore JS, Shaw C, Shaw E, *et al.* Melanoma in horses: Current perspectives. *Equine Vet Educ* 2013; **25**: 144–151.
105. Seltenhammer MH, Simhofer H, Scherzer S, *et al.* Equine melanoma in a population of 296 grey Lipizzaner horses. *Equine Vet J* 2003; **35**: 153–157.
106. Campagne C, Julé S, Bernex F, *et al.* RACK1, a clue to the diagnosis of cutaneous melanomas in horses. *BMC Vet Res* 2012; **8**: 95.
107. Rosengren Pielberg G, Golovko A, Sundström E, *et al.* A cis-acting regulatory mutation causes premature hair graying and susceptibility to melanoma in the horse. *Nat Genet* 2008; **40**: 1004–1009.
108. Sundström E, Komisarczuk AZ, Jiang L, *et al.* Identification of a melanocytespecific, microphthalmia-associated transcription factor-dependent regulatory element in the intronic duplication causing hair graying and melanoma in horses. *Pigment Cell Melanoma Res* 2012; **25**: 28–36.
109. Jiang L, Campagne C, Sundström E, *et al.* Constitutive activation of the ERK pathway in melanoma and skin melanocytes in Grey horses. *BMC Cancer* 2014; **14**: 857.
110. Kansal RG, McCravy MS, Basham JH, *et al.* Inhibition of melanocortin 1 receptor slows melanoma growth, reduces tumor heterogeneity and increases survival. *Oncotarget* 2016; **7**: 26331–26345.

111. Fleury C, Berard F, Balme B *et al.* The study of cutaneous melanomas in Camargue-type gray-skinned horses (1): Clinical-pathological characterization. *Pigment Cell Res* 2000; **13**: 39–46.
112. Wong K, van der Weyden L, Schott CR, *et al.* Cross-species genomic landscape comparison of human mucosal melanoma with canine oral and equine melanoma. *Nat Commun* 2019; **10**: 353.
113. Hayward NK, Wilmott JS, Waddell N, *et al.* Whole-genome landscapes of major melanoma subtypes. *Nature* 2017; **545**: 175–180.
114. Bax MJ, Brown MD, Rothberg PG, *et al.* Pigmented epithelioid melanocytoma (animal-type melanoma): an institutional experience. *J Am Acad Dermatol* 2017; **77**: 328–332.
115. Patnaik AK, Liu SK, Hurvitz AI, *et al.* Nonhematopoietic neoplasms in cats. *J Natl Cancer Inst* 1975; **54**: 855–860.
116. Patnaik AK, Mooney S. Feline melanoma: a comparative study of ocular, oral, and dermal neoplasms. *Vet Pathol* 1988; **25**: 105–112.
117. Bourguet A, Piccicuto V, Donzel E, *et al.* A case of primary choroidal malignant melanoma in a cat. *Vet Ophthalmol* 2015; **18**: 345–349.
118. Stei MM, Loeffler KU, Holz FG, *et al.* Animal Models of Uveal Melanoma: Methods, Applicability, and Limitations. *Biomed Res Int* 2016; **2016**: 4521807.
119. Gillard M, Cadieu E, De Brito C, *et al.* Naturally occurring melanomas in dogs as models for non-UV pathways of human melanomas. *Pigment Cell Melanoma Res* 2014; **27**: 90–102.
120. Ramos-Vara JA, Beissenherz ME, Miller MA, *et al.* Retrospective study of 338 canine oral melanomas with clinical, histologic, and immunohistochemical review of 129 cases. *Vet Pathol* 2000; **37**: 597–608.
121. Prouteau A, Andre C. Canine Melanomas as Models for Human Melanomas: Clinical, Histological, and Genetic Comparison. *Genes* 2019; **10**: 501.

122. Hernandez B, Adissu HA, Wei BR, *et al.* Naturally Occurring Canine Melanoma as a Predictive Comparative Oncology Model for Human Mucosal and Other Triple Wild-Type Melanomas. *Int J Mol Sci* 2018; **19**: 394.
123. Bergman P, Wolchok J. Of Mice and Men (and Dogs): development of a xenogeneic DNA vaccine for canine oral malignant melanoma. *Cancer Ther* 2008; **6**: 817–826.
124. D'Angelo SP, Larkin J, Sosman JA, *et al.* Efficacy and Safety of Nivolumab Alone or in Combination With Ipilimumab in Patients With Mucosal Melanoma: A Pooled Analysis. *J Clin Oncol* 2017; **35**: 226–235.
125. Maekawa N, Konnai S, Takagi S, *et al.* A canine chimeric monoclonal antibody targeting PD-L1 and its clinical efficacy in canine oral malignant melanoma or undifferentiated sarcoma. *Sci Rep* 2017; **7**:8951.
126. Alexander AN, Huelsmeyer MK, Mitzey A, *et al.* Development of an allogeneic whole-cell tumor vaccine expressing xenogeneic gp100 and its implementation in a phase II clinical trial in canine patients with malignant melanoma. *Cancer Immunol Immunother* 2006; **55**: 433–442.
127. Bianco SR, Sun J, Fosmire SP, *et al.* Enhancing antimelanoma immune responses through apoptosis. *Cancer Gene Ther* 2003; **10**: 726–736.
128. Verganti S, Berlato D, Blackwood L, *et al.* Use of Oncept melanoma vaccine in 69 canine oral malignant melanomas in the UK. *J Small Anim Pract* 2017; **58**: 10–16.
129. Ottnod JM, Smedley RC, Walshaw R, *et al.* A retrospective analysis of the efficacy of Oncept vaccine for the adjunct treatment of canine oral malignant melanoma. *Vet Comp Oncol* 2013; **11**: 219–229.
130. Simpson RM, Bastian BC, Michael HT, *et al.* Sporadic naturally occurring melanoma in dogs as a preclinical model for human melanoma. *Pigment Cell Melanoma Res* 2014; **27**: 37–47.

131. Hendricks WPD, Zismann V, Sivaprakasam K, *et al.* Somatic inactivating PTPRJ mutations and dysregulated pathways identified in canine malignant melanoma by integrated comparative genomic analysis. *PLoS Genet* 2018; **14**: e1007589.
132. Newell F, Kong Y, Wilmott JS, *et al.* Whole-genome landscape of mucosal melanoma reveals diverse drivers and therapeutic targets. *Nat Commun* 2019; **10**: 3163.
133. Fowles JS, Denton CL, Gustafson DL. Comparative analysis of MAPK and PI3K/AKT pathway activation and inhibition in human and canine melanoma. *Vet Comp Oncol* 2015; **13**: 288–304.
134. Küsters-Vandeveldde HVN, Germans MR, Rabbie R, *et al.* Whole-exome sequencing of a meningeal melanocytic tumour reveals activating CYSLTR2 and EIF1AX hotspot mutations and similarities to uveal melanoma. *Brain Tumor Pathol* 2018; **35**: 127–130.
135. Hitte C, Le Béguec C, Cadieu E, *et al.* Genome-Wide Analysis of Long Non-Coding RNA Profiles in Canine Oral Melanomas. *Genes (Basel)* 2019; **10**: 477.
136. Smith SH, Goldschmidt MH, McManus PM. A comparative review of melanocytic neoplasms. *Vet Pathol* 2002; **39**: 651–678.
137. Dobson JM. Breed-predispositions to cancer in pedigree dogs. *ISRN Vet Sci* 2013; **2013**: 941275.
138. Hendricks WPD, Zismann V, Sivaprakasam K, *et al.* Somatic inactivating PTPRJ mutations and dysregulated pathways identified in canine malignant melanoma by integrated comparative genomic analysis. *PLoS Genet* 2018; **14**: e1007589.
139. Giuliano EA, Chappell R, Fischer B. A matched observational study of canine survival with primary intraocular melanocytic neoplasia. *Vet Ophthalmol* 1999; **2**: 185–190.
140. Onken MD, Worley LA, Tuscan MD, *et al.* An accurate, clinically feasible multi-gene expression assay for predicting metastasis in uveal melanoma. *J Mol Diagn* 2010; **12**: 461–468.

141. Malho P, Dunn K, Donaldson D, *et al.* Investigation of prognostic indicators for human uveal melanoma as biomarkers of canine uveal melanoma metastasis. *J Small Anim Pract* 2013; **54**: 584–593.
142. Henry CJ, Brewer WG, Whitley EM, *et al.* Canine digital tumors: A veterinary cooperative oncology group retrospective study of 64 dogs. *J Vet Intern Med* 2005; **19**: 720–724.
143. Wobeser BK, Kidney BA, Powers BE, *et al.* Diagnoses and clinical outcomes associated with surgically amputated canine digits submitted to multiple veterinary diagnostic laboratories. *Vet Pathol* 2007; **44**: 355–361.
144. Kim DY, Royal AB, Villamil JA. Disseminated melanoma in a dog with involvement of leptomeninges and bone marrow. *Vet Pathol* 2009; **46**: 80–83.
145. Snyder JM, Lipitz L, Skorupski KA, *et al.* Secondary intracranial neoplasia in the dog: 177 cases (1986-2003). *J Vet Intern Med* 2008; **22**: 172–177.
146. Song RB, Vite CH, Bradley CW, *et al.* Postmortem evaluation of 435 cases of intracranial neoplasia in dogs and relationship of neoplasm with breed, age, and body weight. *J Vet Intern Med* 2013; **27**: 1143–1152.

Table 1. Summary of key genetic alterations of the different melanoma classes.

Sun status	High CSD	Low CSD	Sun protected areas			
Tissue	Cutaneous		Acral	Mucosal	Uveal	CNS
Oncogene activation	<i>NRAS</i> <i>BRAF</i> ¹ <i>KIT</i>	<i>BRAF</i> <i>NRAS</i>	<i>BRAF</i> <i>NRAS</i> <i>GNAQ</i> <i>KIT</i> <i>CCND1</i> (<i>amp</i>)	<i>NRAS</i> <i>KIT</i> <i>BRAF</i> <i>SF3B1</i> <i>CCND1</i> (<i>amp</i>)	<i>GNAQ</i> <i>GNA11</i> <i>CYSLTR2</i> <i>PLCB4</i> <i>SF3B1</i>	<i>GNAQ</i> <i>GNA11</i> <i>CYSLTR2</i> <i>PLCB4</i> <i>SF3B1</i> <i>NRAS</i> ²
Tumour suppressor gene loss	<i>NF1</i> <i>TP53</i> <i>CDKN2A</i> <i>PTEN</i>	<i>TP53</i> <i>CDKN2A</i> <i>PTEN</i>	<i>NF1</i> <i>CDKN2A</i>	<i>NF1</i> <i>PTEN</i> <i>TP53</i> <i>CDK4</i>	<i>BAP1</i>	<i>BAP1</i>
Others	<i>TERT</i> (<i>prom</i>)	<i>TERT</i> (<i>prom</i>)	<i>TERT</i> (<i>amp</i>)	<i>TERT</i> (<i>amp</i>)	<i>EIF1AX</i>	<i>EIF1AX</i>

Abbreviations: Amp, amplification; CNS, central nervous system; CSD, cumulative sun damage; Prom; promoter mutations. ¹The *BRAF* mutation is not the typical *V600E* variant. ²There is a strong link to *NRAS* mutations in cases of childhood melanomas and neurocutaneous melanocytosis.

Table 2. Some examples of the wide range of species in which spontaneously occurring melanoma or melanocytoma has been reported.

Animal	Clinical presentation	Diagnosis and histopathology report	Outcome	Ref
Nurse shark (<i>Ginglymostoma cirratum</i>)	A 5.5-yr history of a 6-cm black, raised nodular skin lesion located on the right side of the proximal tail.	Cutaneous melanoma. The biopsy comprised interlacing streams of neoplastic spindle cells with scant eosinophilic cytoplasm and vesicular nuclei with prominent nucleoli. A few of the cells contained melanin. No vascular invasion was noted.	Euthanized for systemic illness approximately 4.5 months after diagnosis (although no evidence of metastasis found on histopathologic evaluation of the skin and viscera)	31
Haller's round ray (<i>Urobatis halleri</i>)	Presented with multiple black raised nodular masses on the dorsal surface.	Cutaneous malignant melanoma. The masses were composed of proliferative sheets of melanocytes exhibiting mild anisocytosis and anisokaryosis.	Approximately 2 months following the biopsy, the ray became acutely anorexic and was found dead.	32
Coral trout (<i>Plectropomus leopardus</i>)	20/136 of line caught coral trout from two locations in the Great Barrier Reef Marine Park (in 2010-12) showed a dark growth lesion covering <10% to the entire body.	Melanosis and melanoma. The lesions showed a tumorous appearance of disorganised pleomorphic cells containing melanosomes, consistent with melanophorous-macromelanophorous polymorphic melanoma (MMPM).	N/A.	33
Florida pine snake (<i>Elaphe obsoleta rossalleni</i>)	Presented with a black dermal tumour, dorso-laterally in the posterior third of the body.	Malignant melanoma. The melanotic cells were polyhedral and uniformly round or nearly round. The vesicular nuclei occupied most of the cellular space and the nucleoli were prominent. The scanty amount of protoplasm was heavily impregnated with melanin granules. Widespread metastasis.	Two months after tumour excision, further small black tumours appeared in various parts of the skin and the general condition of the snake began to deteriorate; it died soon after.	34
Mandarin duck (<i>Aix galericulata</i>)	Presented with several small multinodular masses on the dorsal surface of the upper bill.	Metastatic malignant Melanoma. The tissue consisted of multiple nests and lobules of neoplastic melanocytes (with prominent areas of necrosis). The lymphoid tissues contained nests and individual neoplastic melanocytes.	Two months later, the tumour had enlarged considerably. The duck developed severe dyspnea and was euthanatized.	35
Macaroni penguin (<i>Eudyptes chrysolophus</i>)	Presented with a caseous and necrotic mass that engulfed the upper beak and extended into the rostra sinuses.	Malignant melanoma. The tissue contained an infiltrative, deeply pigmented melanocytic neoplasm that extended from superficial to deep dermal regions and had effaced normal tissue architecture.	The penguin exhibited dyspnea as a result of invasion of the tumour into the rostral sinuses, and was immediately euthanised.	36
Rabbit (<i>Oryctolagus cuniculus</i>)	Presented with an irregular black mass (4 × 3 × 2 cm) on the skin of scrotum, with ulceration and bleeding.	Cutaneous malignant melanoma. The tumour cells were positive for a number of melanocyte markers (HMB-45, PNL2, Melan A, and S100) and Ki-67.	The perioperative period concluded with no problems, however, despite subsequent clinical progress, the rabbit died after two weeks due to metastasis.	37

Pygora goat <i>(Capra aegagrus hircus)</i>	Presented with a recurrent ulcerated, black-pigmented, 2.5 cm mass at the base of the left horn.	Malignant melanoma. The masses consisted of moderately pleomorphic, polyhedral to spindle cells containing variable amounts of dark brown intracytoplasmic pigment granules. Multiple black foci of metastasis observed in the liver.	While restraining the goat for physical examination the left horn broke off at the base with minimal force. Due to the poor condition of the goat it was immediately euthanized.	38
Huacaya alpaca <i>(Vicugna pacos)</i>	Presented with a chronic, non-healing wound involving the left external nostril.	Metastatic mucocutaneous melanoma. Malignant melanoma was diagnosed by histology of biopsy specimens. At post-mortem, numerous masses present throughout the body were composed of spindloid to polygonal cells with indistinct cell borders and moderate amounts of cytoplasm containing abundant melanin.	The alpaca was euthanized 10 days after the diagnosis on the basis of the poor prognosis and rapid clinical deterioration.	39
Pygmy hippopotamus <i>(Choeropsis liberiensis)</i>	Presented with multiple raised and pigmented skin masses.	Dermal malignant melanoma and concurrent melanocytoma. Initial impression smears of one ulcerated lesion were consistent with inflammation however, subsequent histopathological findings from a skin biopsy revealed an underlying malignant melanoma.	There was no sign of recurrence 34 months post-surgery (no lymph node involvement).	40
African lion <i>(Panthera leo)</i>	Presented with a 4 month history of left maxillary lip swelling. Examination showed a pigmented mass at the level of the left maxillary canine tooth.	Dermal malignant melanoma. Histopathologic evaluation of the biopsies revealed a malignant dermal melanoma with no evidence of metastasis.	The lion received radiotherapy and immunotherapy treatments.	41
Aberdeen Angus cow <i>(Bos taurus)</i>	Presented with a large, pedunculated cutaneous mass protruding from the left flank fold and an enlarged left pre-femoral lymph node.	Congenital amelanotic melanoma with nodal metastasis. Histologic examination revealed a homogeneous population of neoplastic cells staining positively for S100 and melan A.	Two months later, the calf became acutely recumbent and was euthanized after clinical examination revealed widespread metastasis.	42
Cynomolgus monkey <i>(Macaca fascicularis)</i>	A pigmented raised mass (2 cm diameter) at the dorsal aspect of the neck.	Cutaneous melanocytoma. Histologically, the mass was composed of poorly demarcated, heavily pigmented melanocytes diffusely infiltrating the dermis. The neoplastic cells were fairly uniform round to polygonal in shape and arranged in packets, supported by a fibrovascular stroma. The nuclei were mostly round and lacked nuclear atypia.	Not reported.	43

Table 3. Key features of each animal as a 'model' of melanoma in humans.

Animal	Key features as a model
Platyfish/ Swordtail	<ul style="list-style-type: none"> • Melanoma arises from the EGFR pathway (uses Ras/Raf/MAPK signalling) • Melanoma incidence controlled by a pigment-cell-specific oncogene locus and a tumour suppressor locus • Model of UV radiation-induced melanoma
Gray short-tailed opossum	<ul style="list-style-type: none"> • Model of UV radiation-induced melanoma • Model of sunscreen-mediated melanoma prevention
Miniature pig	<ul style="list-style-type: none"> • Opportunity to study the entire spectrum of melanoma formation, from benign lesions to malignant transformation to metastatic spread • Model of spontaneous regression of melanomas
Horse	<ul style="list-style-type: none"> • Melanomas under the tail and in the peri-anal region, lips and eyelids are models of mucosal melanoma • Driver genes in these mucosal melanomas: <i>NRAS</i>, <i>TP53</i>, <i>PTEN</i>, <i>KIT</i> and <i>BRAF</i>
Cat	<ul style="list-style-type: none"> • Melanomas of the uvea are a model of uveal melanoma
Dog	<ul style="list-style-type: none"> • Melanomas of the oral cavity are a model of mucosal melanoma • Driver genes in these mucosal melanomas: <i>NRAS</i>, <i>TP53</i>, <i>PTEN</i>, <i>NF1</i>, <i>KRAS</i>, and <i>BRCA2</i> • MAPK and PI3K/AKT pathways involved in mucosal melanoma formation and progression • Melanomas of the footpad or nail bed are models of acral melanoma • Clinical trials are performed in dogs with melanoma

Figure legends

Figure 1. The two major signaling pathways implicated in melanoma. Commonly mutated genes underlined. Receptors are shown in blue, proteins in the PI3K/AKT pathway are in yellow, proteins in the MAPK/ERK pathway are in green, proteins in the GNAQ/GNA11 pathway are in pink and proteins that have inhibitory roles in these pathways are in red.

Figure 2. Metastatic melanophoroma in a red-eared slider turtle. (A, B) Multifocal malignant melanophoromas on the shell (circle). (C, D) The lesions invaded through the shell (circle) into the peritoneal cavity (arrows). The photographs were kindly provided by Luke Haydock, Department of Pathobiology, University of Guelph, Guelph, Ontario, Canada.

Figure 3. Melanoma in a Xiphophorus. (A) Macroscopic photo of a platyfish/swordtail backcross hybrid (Tu+/-, R-/-) with a large pigmented melanoma in the posterior and on the pectoral fin (arrows). (B-D) Images of H&E-stained sections of a large exophytic melanoma that covers the surface of the skin (circle), while heavily pigmented melanoma cells can be seen infiltrating the muscle (arrows) (25x, 100x and 200x magnification, respectively).

Figure 4. Melanomas in Sinclair miniature pigs. (A) A cutaneous melanoma on the flank (arrow). (B) A regressed cutaneous melanoma showing areas of skin depigmentation (arrow). (C) Large cutaneous melanomas on the belly and inner hind limb (arrows). (D) Hepatic metastases from a cutaneous melanoma (arrows). The photographs were kindly provided by Derek Brocksmith, Sinclair BioResources, Missouri, USA.

Figure 5. Metastatic melanoma in a horse. (A) Multiple melanomas located under the tail base (circle) that metastasized to the lungs and mediastinum around the heart. Shown are thoracic metastases (B; arrows) and metastases on the pleural surface of the ribs (C; arrows). The

1 photographs were kindly provided by Laura Bassel, Department of Pathobiology, University of
2
3
4 Guelph, Guelph, Ontario, Canada.
5
6
7

8 **Figure 6.** Uveal melanoma in a cat. A diffuse iris melanoma (arrow). The photograph was
9
10 kindly provided by Chantale Pinard, Department of Clinical Studies, University of Guelph,
11
12 Guelph, Ontario, Canada.
13
14
15
16
17
18
19

20 **Figure 7.** Oral and uveal melanomas in dogs. (A) A primary melanoma in the soft palate (inset),
21
22 that invaded into oropharynx (arrow). The photographs were kindly provided by Laura Bassel,
23
24 Department of Pathobiology, University of Guelph, Guelph, Ontario, Canada. (B) A benign
25
26 uveal melanoma (melanocytoma) in the iris (arrow). The photograph was kindly provided by
27
28 Chantale Pinard, Department of Clinical Studies, University of Guelph, Guelph, Ontario,
29
30 Canada.
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 1. The two major signaling pathways implicated in melanoma. Commonly mutated genes underlined. Receptors are shown in blue, proteins in the PI3K/AKT pathway are in yellow, proteins in the MAPK/ERK pathway are in green, proteins in the GNAQ/GNA11 pathway are in pink and proteins that have inhibitory roles in these pathways are in red.

338x190mm (54 x 54 DPI)




Figure 2. Metastatic melanophoroma in a red-eared slider turtle. (A, B) Multifocal malignant melanophoromas on the shell (circle). (C, D) The lesions invaded through the shell (circle) into the peritoneal cavity (arrows). The photographs were kindly provided by Luke Haydock, Department of Pathobiology, University of Guelph, Guelph, Ontario, Canada.

338x190mm (54 x 54 DPI)




Figure 3. Melanoma in a Xiphophorus. (A) Macroscopic photo of a platyfish/swordtail backcross hybrid (Tu+/-, R-/-) with a large pigmented melanoma in the posterior and on the pectoral fin (arrows). (B-D) Images of H&E-stained sections of a large exophytic melanoma that covers the surface of the skin (circle), while heavily pigmented melanoma cells can be seen infiltrating the muscle (arrows) (25x, 100x and 200x magnification, respectively).

338x190mm (54 x 54 DPI)




Figure 4. Melanomas in Sinclair miniature pigs. (A) A cutaneous melanoma on the flank (arrow). (B) A regressed cutaneous melanoma showing areas of skin depigmentation (arrow). (C) Large cutaneous melanomas on the belly and inner hind limb (arrows). (D) Hepatic metastases from a cutaneous melanoma (arrows). The photographs were kindly provided by Derek Brocksmith, Sinclair BioResources, Missouri, USA.

338x190mm (54 x 54 DPI)




Figure 5. Metastatic melanoma in a horse. (A) Multiple melanomas located under the tail base (circle) that metastasized to the lungs and mediastinum around the heart. Shown are thoracic metastases (B; arrows) and metastases on the pleural surface of the ribs (C; arrows). The photographs were kindly provided by Laura Bassel, Department of Pathobiology, University of Guelph, Guelph, Ontario, Canada.

338x190mm (54 x 54 DPI)




Figure 6. Uveal melanoma in a cat. A diffuse iris melanoma (arrow). The photograph was kindly provided by Chantale Pinard, Department of Clinical Studies, University of Guelph, Guelph, Ontario, Canada.

338x190mm (54 x 54 DPI)




Figure 7. Oral and uveal melanomas in dogs. (A) A primary melanoma in the soft palate (inset), that invaded into oropharynx (arrow). The photographs were kindly provided by Laura Bassel, Department of Pathobiology, University of Guelph, Guelph, Ontario, Canada. (B) A benign uveal melanoma (melanocytoma) in the iris (arrow). The photograph was kindly provided by Chantale Pinard, Department of Clinical Studies, University of Guelph, Guelph, Ontario, Canada.

338x190mm (54 x 54 DPI)